

Leitgraptoliten	Mächtigkeit (m)	Stratigraphie	
Monograptus hercynicus	8-12	Devon Lochkow	
Monograptus praehercynicus	1,5-3		
Monograptus uniformis	3,5-5		
Monogr. (Pristiograptus) transgrediens	10-25	Budenium	
Monogr. (Pristiogr.) dubius thuringicus			
Monogr. (Saetograptus) fritschi linearis	5-15		
Monogr. (Saetograptus) leintwardinensis			
Monograptus (Saetograptus) chimera	3-10		
Monograptus scanicus			
Monograptus (Pristiograptus) nilssoni	1,8		
Monograptus (Pristiograptus) vulgaris	1,5		
Mongraptus deubeli	0,3		
Monograptus testis	2-3		Wenlock
Cyrtograptus radians	1,2		
(Cyrtograptus ellesae)	1-3		
Monograptus flexilis			
(Cyrtograptus rigidus)			
Monograptus riccartonensis	0,7		
Cyrtograptus murchisoni			
Monograptus (Spirograptus) spiralis	3,8	Llandovery	
Monograptus (Monoclimacis) crenulata	1,2		
Monogr. (Monoclimacis) griestonensies	1,0		
Monograptus (Spirograptus) turriculatus	1,0		
Rastrites linnaei	3-5		
Monograptus sedgwicki	4,5-5		
Rastrites peregrinus	1,7		
Monograptus (Demirastrites) convolutus	1,5-2		
Monograptus (Pristiograptus) gregarius	1,5		
Monograptus (Pristiograptus) cyphus	0,8-1		
Orthograptus vesiculosus	0,4-1		
Akidograptus acuminatus	0,4-5		
Akidograptus ascensus	0,4		
Glyptograptus persculptus	?		
Dicellograptus anceps	?	Ordovizium	

www.regionalgeologie-ost.de

Computergrafik: D. FRANKE

Abb. 34.3.1 Graptolithenzonen-Abfolge im Präperm (Gräfenwarth-Gruppe) des Thüringischen Schiefergebirges

(nach G. SCHLEGEL (1998a))